

surgery on the cervix or uterus including cesarean section, overdistention of the uterus, grand multiparity, or past history of uterine sepsis or of traumatic delivery. Because of the variability of the combinations of factors which may be present in the conditions listed above, the definition of "unusual circumstances" must be left to the judgment of the physician. The decision can be made only by carefully weighing the potential benefits which oxytocin can provide in a given case against rare but definite potential for the drug to produce hypertonicity or tetanic spasm.

4. Maternal deaths due to hypertensive episodes, subarachnoid hemorrhage, rupture of the uterus, and fetal deaths due to various causes have been reported associated with the use of parenteral oxytocic drugs for induction of labor or for augmentation in the first and second stages of labor.
5. Oxytocin has been shown to have an intrinsic antidiuretic effect, acting to increase water reabsorption from the glomerular filtrate. Consideration should, therefore, be given to the possibility of water intoxication, particularly when oxytocin is administered continuously by infusion and the patient is receiving fluids by mouth.
6. When oxytocin is used for induction or reinforcement of already existent labor, patients should be carefully selected. Pelvic adequacy must be considered and maternal and fetal conditions evaluated before use of the drug.

Drug Interactions

Severe hypertension has been reported when oxytocin was given three to four hours following prophylactic administration of a vasoconstrictor in conjunction with caudal block anesthesia. Cyclopropane anesthesia may modify oxytocin's cardiovascular effects, so as to produce unexpected results such as hypotension. Maternal sinus bradycardia with abnormal atrioventricular rhythms has also been noted when oxytocin was used concomitantly with cyclopropane anesthesia.

Carcinogenesis, Mutagenesis, Impairment of Fertility

There are no animal or human studies on the carcinogenicity and mutagenicity of this drug, nor is there any information on its effect on fertility.

Pregnancy

Teratogenic Effects

Animal reproduction studies have not been conducted with oxytocin. There are no known indications for use in the first trimester of pregnancy other than in relation to spontaneous or induced abortion. Based on the wide experience with this drug and its chemical structure and pharmacological properties, it would not be expected to present a risk of fetal abnormalities when used as indicated.

Nonteratogenic Effects

See ADVERSE REACTIONS in the fetus or neonate.

Labor and Delivery

See INDICATIONS AND USAGE section.

ADVERSE REACTIONS

The following adverse reactions have been reported in the mother:

Anaphylactic reaction	Premature ventricular contractions
Postpartum hemorrhage	Pelvic hematoma
Cardiac arrhythmia	Subarachnoid hemorrhage
Fatal afibrinogenemia	Hypertensive episodes
Nausea	Rupture of the uterus
Vomiting	

Excessive dosage or hypersensitivity to the drug may result in uterine hypertonicity, spasm, tetanic contraction, or rupture of the uterus.

The possibility of increased blood loss and afibrinogenemia should be kept in mind when administering the drug.

Severe water intoxication with convulsions and coma has occurred, associated with a slow oxytocin infusion over a 24-hour period. Maternal death due to oxytocin-induced water intoxication has been reported.

The following adverse reactions have been reported in the fetus or neonate:

Due to induced uterine motility:	Due to use of oxytocin in the mother:
Bradycardia	Low Apgar scores at five minutes
Premature ventricular contractions and other arrhythmias	Neonatal jaundice
Permanent CNS or brain damage	Neonatal retinal hemorrhage
Fetal death	
Neonatal seizures have been reported with the use of Pitocin.	

OVERDOSAGE

Overdosage with oxytocin depends essentially on uterine hyperactivity whether or not due to hypersensitivity to this agent. Hyperstimulation with strong (hypertonic) or prolonged (tetanic) contractions, or a resting tone of 15 to 20 mm H₂O or more between contractions can lead to tumultuous labor, uterine rupture, cervical and vaginal lacerations, postpartum hemorrhage, uteroplacental hypoperfusion, and variable deceleration of fetal heart, fetal hypoxia, hypercapnia, perinatal hepatic necrosis or death. Water intoxication with convulsions, which is caused by the inherent antidiuretic effect of oxytocin, is a serious complication that may occur if large doses (40 to 50 milliunits/minute) are infused for long periods. Management consists of immediate discontinuation of oxytocin and symptomatic and supportive therapy.

DOSAGE AND ADMINISTRATION

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration whenever solution and container permit.

The dosage of oxytocin is determined by the uterine response and must therefore be individualized and initiated at a very low level. The following dosage information is based upon various regimens and indications in general use.

A. Induction or Stimulation of Labor

Intravenous infusion (drip method) is the only acceptable method of parenteral administration of Pitocin for the induction or stimulation of labor. Accurate control of the rate of infusion is essential and is best accomplished by an infusion pump. It is convenient to piggyback the Pitocin infusion on a physiologic electrolyte solution, permitting the Pitocin infusion to be stopped abruptly without interrupting the electrolyte infusion. This is done in the following way.

1. Preparation

- a. The standard solution for infusion of Pitocin is prepared by adding the contents of one 1-mL vial containing 10 units of oxytocin to 1000 mL of 0.9% aqueous sodium chloride or Ringer's lactate. The combined solution containing 10 milliunits (mU) of oxytocin/mL is rotated in the infusion bottle for thorough mixing.
- b. Establish the infusion with a separate bottle of physiologic electrolyte solution not containing Pitocin.

- c. Attach (piggyback) the Pitocin-containing bottle with the infusion pump to the infusion line as close to the infusion site as possible.

2. Administration

The initial dose should be 0.5–1 mU/min (equal to 3–6 mL of the dilute oxytocin solution per hour). At 30–60 minute intervals the dose should be gradually increased in increments of 1–2 mU/min until the desired contraction pattern has been established. Once the desired frequency of contractions has been reached and labor has progressed to 5–6 cm dilation, the dose may be reduced by similar increments.

Studies of the concentrations of oxytocin in the maternal plasma during Pitocin infusion have shown that infusion rates up to 6 mU/min give the same oxytocin levels that are found in spontaneous labor. At term, higher infusion rates should be given with great care, and rates exceeding 9–10 mU/min are rarely required. Before term, when the sensitivity of the uterus is lower because of a lower concentration of oxytocin receptors, a higher infusion rate may be required.

3. Monitoring

- a. Electronically monitor the uterine activity and the fetal heart rate throughout the infusion of Pitocin. Attention should be given to toonus, amplitude and frequency of contractions, and to the fetal heart rate in relation to uterine contractions. If uterine contractions become too powerful, the infusion can be abruptly stopped, and oxytocic stimulation of the uterine musculature will soon wane (see **PRECAUTIONS** section).
- b. Discontinue the infusion of Pitocin immediately in the event of uterine hyperactivity and/or fetal distress. Administer oxygen to the mother, who preferably should be put in a lateral position. The condition of mother and fetus should immediately be evaluated by the responsible physician and appropriate steps taken.

B. Control of Postpartum Uterine Bleeding

1. Intravenous infusion (drip method). If the patient has an intravenous infusion running, 10 to 40 units of oxytocin may be added to the bottle, depending on the amount of electrolyte or dextrose solution remaining (maximum 40 units to 1000 mL). Adjust the infusion rate to sustain uterine contraction and control uterine atony.
2. Intramuscular administration. (One mL) Ten (10) units of Pitocin can be given after the delivery of the placenta.

C. Treatment of Incomplete, Inevitable, or Elective Abortion

Intravenous infusion of 10 units of Pitocin added to 500 mL of a physiologic saline solution or 5% dextrose-in-water solution may help the uterus contract after a suction or sharp curettage for an incomplete, inevitable, or elective abortion.

Subsequent to intra-amniotic injection of hypertonic saline, prostaglandins, urea, etc., for midtrimester elective abortion, the injection-to-abortion time may be shortened by infusion of Pitocin at the rate of 10 to 20 milliunits (20 to 40 drops) per minute. The total dose should not exceed 30 units in a 12-hour period due to the risk of water intoxication.

HOW SUPPLIED

Pitocin (Oxytocin Injection, USP) Synthetic is available as follows: NDC 61570-416-05 Packages of twenty-five oversized 1-mL Steri-Vials®, each containing 10 units of oxytocin.

STORAGE

Store at 2°–8°C (36°–46°F). May be held at 15°–25°C (59°–77°F) for up to 30 days. Discard after holding at 15°–25°C (59°–77°F).

REFERENCES

1. Seitchik J, Castillo M: Oxytocin augmentation of dysfunctional labor. I. Clinical data. *Am J Obstet Gynecol* 1982; 144:899–905.
2. Seitchik J, Castillo M: Oxytocin augmentation of dysfunctional labor. II. Multiparous patients. *Am J Obstet Gynecol* 1983; 145:777–780.
3. Fuchs A, Goeschen K, Husslein P, et al: Oxytocin and the initiation of human parturition. III. Plasma concentrations of oxytocin and 13, 14-dihydro-15-keto-prostaglandin F2a in spontaneous and oxytocin-induced labor at term. *Am J Obstet Gynecol* 1983; 145:497–502.
4. Seitchik J, Amico J, et al: Oxytocin augmentation of dysfunctional labor. IV. Oxytocin pharmacokinetics. *Am J Obstet Gynecol* 1984; 150:225–228.
5. American College of Obstetricians and Gynecologists: ACOG Technical Bulletin Number 110—November 1987: Induction and augmentation of labor.

Rx only.

Prescribing Information as of November 2004.



Distributed by:
Monarch Pharmaceuticals, Inc., Bristol, TN 37620
Manufactured by:
Parkedale Pharmaceuticals, Inc., Rochester, MI 48307